

# ON THE ASPHERICITY OF KNOT COMPLEMENTS

VO THANH LIEM AND GERARD A. VENEMA

ABSTRACT. Two examples of topological embeddings of  $S^2$  in  $S^4$  are constructed. The first has the unusual property that the fundamental group of the complement is isomorphic to the integers while the second homotopy group of the complement is nontrivial. The second example is a non-locally flat embedding whose complement exhibits this property locally.

Two theorems are proved. The first answers the question of just when good  $\pi_1$  implies the vanishing of the higher homotopy groups for knot complements in  $S^4$ . The second theorem characterizes local flatness for 2-spheres in  $S^4$  in terms of a local  $\pi_1$  condition.

## 0. INTRODUCTION

In this paper we construct two examples of topological embeddings of  $S^2$  into  $S^4$  whose complements have unusual homotopy properties. In order to explain the context in which we are working, we begin with a question whose answer is well known. The question is this: If  $\Sigma$  is a (topologically) locally flat  $(n - 2)$ -sphere in  $S^n$  such that  $\pi_1(S^n - \Sigma) \cong \mathbb{Z}$ , then do the higher homotopy groups of  $S^n - \Sigma$  necessarily vanish? (In case they do,  $S^n - \Sigma$  is said to be *aspherical* and will have the homotopy type of  $S^1$ .) The answer to this question is “yes” in case  $n \leq 4$  and “no” in case  $n \geq 5$ . For  $n = 3$  this follows from the sphere theorem of Papakyriakopoulos [14]. In case  $n = 4$ , there are two quite different arguments that can be used. Both proofs begin by noting that, since  $\Sigma$  is locally flat, there is a strong deformation retraction of  $S^4 - \Sigma$  to a compact 4-manifold with  $S^2 \times S^1$  boundary and so  $S^4 - \Sigma$  has the homotopy type of a finite complex ([2, Corollary 2] or [19, Corollary 5.3]). The proof that  $S^4 - \Sigma$  is aspherical is then completed either by using Milnor Duality [13] to show that all the homology with  $\mathbb{Z}[\pi_1]$ -coefficients of  $S^4 - \Sigma$  vanishes or, alternatively, by using an argument similar to a proof of Wall—see [8] or [11], for example. On the other hand, for  $n \geq 5$  there exist smooth knots  $\Sigma \subset S^n$  such that  $\pi_1(S^n - \Sigma) \cong \mathbb{Z}$  but  $\pi_2(S^n - \Sigma) \neq 0$ . In fact, it is possible [16] to make the first nontrivial homotopy group appear anywhere in the range 2 through  $[(n - 1)/2]$ .

We address two related questions. The first of these questions is: What happens if we drop the hypothesis that  $\Sigma$  be locally flat? It is well known that the answer remains

---

1991 *Mathematics Subject Classification.* 57Q45, 57N45.

*Key words and phrases.* Knot complement, fundamental group, asphericity, 1-*alg*, local flatness, 2-sphere.

Research of the second author partially supported by National Science Foundation Grant DMS-8900822.

unchanged in case  $n = 3$  and in case  $n \geq 5$  (in fact, the proofs referred to above still work in those cases), but we show by example that the answer does change in dimension 4.

**Example 1.1.** *There exists a topological embedding  $h_1 : S^2 \rightarrow S^4$  such that  $\pi_1(S^4 - h_1(S^2)) \cong \mathbb{Z}$  but  $\pi_2(S^4 - h_1(S^2)) \neq 0$ . Furthermore,  $h_1$  is locally flat except at one point.*

If the non-locally flat point is removed from Example 1.1, an interesting embedding of  $\mathbb{R}^2$  into  $\mathbb{R}^4$  results.

**Example 1.2.** *There exists a proper, locally flat, topological embedding  $h_0 : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  such that  $\pi_1(\mathbb{R}^4 - h_0(\mathbb{R}^2)) \cong \mathbb{Z}$  but  $\pi_2(\mathbb{R}^4 - h_0(\mathbb{R}^2)) \neq 0$ .*

When we drop the local flatness hypothesis, there is also a local version of the question considered in this paper. Before we can state it precisely it is necessary to make some definitions.

*Definitions.* Suppose  $N$  is an  $(n - 2)$ -manifold topologically embedded in the interior of the  $n$ -manifold  $M$  and  $x \in N$ . We say that  $N$  is *locally 1-alg at  $x$*  if for every neighborhood  $U$  of  $x$  in  $M$  there exists a neighborhood  $V$  of  $x$  in  $U$  such that any loop in  $V - N$  which is null-homologous in  $V - N$  is also null-homotopic in  $U - N$ . (The “alg” stands for “abelian local groups.” A duality argument shows that the image of  $\pi_1(V - N)$  in  $\pi_1(U - N)$  is isomorphic to  $\mathbb{Z}$ . Other authors, e.g. [6], use the terminology “ $M - N$  has good local  $\pi_1$  at  $x$ ” to mean the same thing.) We say that  $N$  is *locally homotopically unknotted at  $x$*  if  $N$  is locally 1-alg at  $x$  and if, in addition, for every neighborhood  $U$  of  $x$  in  $M$  there exists a neighborhood  $V$  of  $x$  in  $U$  such that the image of  $\pi_i(V - N)$  in  $\pi_i(U - N)$  is trivial for every  $i \geq 2$ .  $\square$

We can now state the second question considered in this paper: If  $N$  is an  $(n - 2)$ -manifold topologically embedded in the interior of the  $n$ -manifold  $M$  and  $N$  is locally 1-alg at  $x \in N$ , then is  $N$  locally homotopically unknotted at  $x$ ? Again the answer to this question is “yes” in case  $n = 3$  [1] and “no” in high dimensions. In order to construct an example for  $n \geq 6$  we proceed as follows. Let  $\Sigma^{n-3} \subset S^{n-1}$  be a locally flat PL  $(n - 3)$ -sphere such that  $\pi_1(S^{n-1} - \Sigma^{n-3}) \cong \mathbb{Z}$  but  $\pi_2(S^{n-1} - \Sigma^{n-3}) \neq 0$ . Take  $\Sigma'$  to be the suspension of  $\Sigma$  in  $S^n$ . Then  $\Sigma'$  is a PL  $(n - 2)$ -sphere in  $S^n$  which is locally 1-alg at every point but is not locally homotopically unknotted at either suspension point. In §2, below, we also construct an example of a 3-sphere in  $S^5$  which is locally 1-alg but not locally homotopically unknotted; however the example is topological and, in fact, we prove that no such 5-dimensional example can be piecewise linear (as the high dimensional example above is). Our second result shows that the answer to the question is also “no” in dimension 4.

**Example 2.1.** *There exists a topological embedding  $h_2 : S^2 \rightarrow S^4$  and a point  $x \in h_2(S^2)$*

such that  $h_2(S^2)$  is locally 1-*alg* at  $x$  but  $h_2(S^2)$  is not locally homotopically unknotted at  $x$ .

This example is surprising and is the main result of this paper. It had generally been believed that the kind of argument used to show that a locally flat 2-sphere in  $S^4$  which has  $\pi_1$  of the complement equal to  $\mathbb{Z}$  is aspherical could be localized to show that locally 1-*alg* implies locally homotopically unknotted in dimension 4. This is stated explicitly in Lemma 9.3B of [6], for example, and is also implicit in [5, Theorem 10]. Our construction in §1 actually contains a counterexample to the proof of [6, Lemma 9.3B]—see Remark 2, below.

The question of whether locally 1-*alg* implies locally homotopically unknotted is an important question because the proof in [6, §9.3] shows that a surface which is locally homotopically unknotted at every point is locally flat. Thus the real question is whether or not local flatness can be characterized in terms of the local  $\pi_1$  condition or if it is necessary to assume, in addition, that the higher local homotopy groups vanish in order to conclude that the surface is locally flat.

In §3 of this paper we consider the problem of giving necessary and sufficient conditions for the complement of a topological 2-knot in  $S^4$  to be aspherical. Our main result in that section (Theorem 3.1) gives such conditions; it shows that the only way in which a 2-knot complement in  $S^4$  with fundamental group  $\mathbb{Z}$  can fail to be aspherical is if  $\pi_2$  of the complement is nontrivial and that this will only happen if the structure of  $\pi_2$  as a module over  $\mathbb{Z}[\pi_1]$  is quite complicated.

**Theorem 3.1.** *Suppose  $\Sigma$  is a 2-sphere topologically embedded in  $S^4$  and  $\pi_1(S^4 - \Sigma) \cong \mathbb{Z}$ . Then the following are equivalent:*

- (1)  $S^4 - \Sigma$  has the homotopy type of  $S^1$ ,
- (2)  $\pi_2(S^4 - \Sigma) = 0$ ,
- (3)  $\pi_2(S^4 - \Sigma)$  is a submodule of a free  $\mathbb{Z}[\pi_1(S^4 - \Sigma)]$ -module, and
- (4)  $\pi_2(S^4 - \Sigma)$  is finitely generated as a module over  $\mathbb{Z}[\pi_1(S^4 - \Sigma)]$ .

**Corollary 3.2.** *If  $\pi_1(S^4 - \Sigma) \cong \mathbb{Z}$  and  $S^4 - \Sigma$  is finitely dominated, then  $S^4 - \Sigma$  has the homotopy type of  $S^1$ .*

**Corollary 3.3.** *If  $\Sigma$  is a PL (not necessarily locally flat) 2-sphere in  $S^4$  such that  $\pi_1(S^4 - \Sigma) \cong \mathbb{Z}$ , then  $S^4 - \Sigma$  has the homotopy type of  $S^1$ .*

**Corollary 3.4.** *If  $\pi_1(S^4 - \Sigma) \cong \mathbb{Z}$ , and  $\pi_1(\epsilon) \cong \mathbb{Z}$ , where  $\epsilon$  is the end of  $S^4 - \Sigma$ , then  $S^4 - \Sigma$  has the homotopy type of  $S^1$ .*

*Definition.* A compact set  $X$  in the interior of a manifold  $M$  is said to be *globally 1-*alg** if for every neighborhood  $U$  of  $X$  in  $M$  there exists a neighborhood  $V$  of  $X$  in  $U$  such that each loop in  $V - X$  which is null-homologous in  $V - X$  is null-homotopic in  $U - X$ .  $\square$

**Corollary 3.5.** *Let  $\Sigma$  be a 2-sphere topologically embedded in  $S^4$ . If  $\Sigma$  is globally 1-*alg* and  $\pi_1(S^4 - \Sigma) \cong \mathbb{Z}$ , then  $S^4 - \Sigma$  has the homotopy type of  $S^1$ .*

Corollaries 3.4 and 3.5 were first proved in [11]. It should be noted that  $\pi_1(\epsilon) \cong \mathbb{Z}$  is not a necessary condition: Guilbault [7] has given an example of a 2-sphere  $\Sigma \subset S^4$  such that  $S^4 - \Sigma$  has the homotopy type of  $S^1$  even though  $\Sigma$  is not globally 1-*alg*. Corollary 3.3 appears in [17].

Example 2.1 raises the question of whether or not a surface in a 4-manifold that is locally 1-*alg* at *every* point will be locally homotopically unknotted. The example does not answer that question because it contains an infinite sequence of points converging to  $x$  at which the embedding fails to be locally 1-*alg*. As was mentioned above, the question is important because a positive answer would have a taming theorem as a corollary. (The proofs in [6, §9.3] show that an embedding of a surface into a 4-manifold which is locally homotopically unknotted at every point will be locally flat.) We do not answer the question in general, but do give positive answers in several special cases, the most important being the case in which the submanifold is a 2-sphere.

**Theorem 4.1.** *Suppose  $\Sigma$  is a 2-sphere topologically embedded in the interior of the 4-manifold  $M^4$ . If the self intersection number of the homology class represented by  $\Sigma$  is 0 and  $\Sigma$  is locally 1-*alg* at  $x$  for every  $x \in \Sigma$ , then  $\Sigma$  is locally homotopically unknotted.*

By [6, §9.3] we have the following corollary.

**Corollary 4.2.** *If  $\Sigma \subset S^4$  is a 2-sphere which is locally 1-*alg* at  $x$  for every  $x \in \Sigma$ , then  $\Sigma$  is locally flat.*

The conclusion of Theorem 4.1 is local. Since we are often able to take an embedding of some surface other than  $S^2$  and make it agree locally with an embedding of  $S^2$ , we can use Theorem 4.1 to obtain information about embeddings of more general surfaces. For example, we have the following two corollaries.

**Corollary 4.3.** *Suppose  $N$  is a surface topologically embedded in the interior of the 4-manifold  $M$ . If  $N$  is locally 1-*alg* at  $x$  for every  $x \in N$  and  $N$  is locally flat except possibly at a closed 0-dimensional set, then  $N$  is locally flat at every point.*

**Corollary 4.4.** *Suppose  $\Sigma$  is a 2-sphere topologically embedded in the interior of the 4-manifold  $M$ . If  $\Sigma$  is locally 1-*alg* at every point and locally flat at at least one point, then  $\Sigma$  is locally flat at every point.*

## 1. CONSTRUCTION OF THE MAIN EXAMPLE

In this section we construct Example 1.1. The construction proceeds indirectly: we begin by constructing an open subset  $W$  of  $S^4$  having the correct homotopy properties

to be the complement of the example we seek and then show how to move  $W$  a little so that  $\Sigma = S^4 - W$  is a topological 2-sphere. We believe that the example is most easily understood if it is thought of as being analogous to the Whitehead continuum in  $S^3$ . In that case, too, it is easiest to begin by constructing the Whitehead manifold which is the complement in  $S^3$  of the Whitehead continuum. Like the Whitehead manifold, our open manifold  $W$  is described as the union of a nested sequence of compact manifolds  $\{W_n\}$ . This similarity with the Whitehead continuum is not accidental; in fact, we will see that  $\Sigma$  can be obtained by starting with the standard 2-sphere  $S^2$  in  $S^4$  and attaching a copy of the Whitehead continuum to  $S^2$ . Specifically, we find a copy  $\text{Wh}$  of the Whitehead continuum which is cellular in  $S^4$  and intersects  $S^2$  in just one point but is tightly wound around  $S^2$ . Then we construct  $\Sigma$  as follows:

$$\Sigma = (S^2 \vee \text{Wh})/\text{Wh} \subset S^4/\text{Wh} \cong S^4.$$

We begin by fixing some notation. Let  $T$  denote an unknotted solid torus in  $\mathbb{R}^3$ , let  $D$  be a flat disk in the interior of  $T$  and  $C = \partial D$ .




FIGURE 1

We think of  $S^4$  as being  $\mathbb{R}^3 \times (-\infty, \infty) \cup \{\infty\}$  and describe level pictures of the manifolds  $W_n$ . Define  $W_2$  to be a regular neighborhood of  $T \times [-1, 0] \cup C \times [0, 1] \cup D \times \{1\}$  in  $S^4$ . Notice that  $W_2$  collapses to a copy of  $S^1 \vee S^2$ . We next define  $W_1 \subset \text{Int } W_2$ . Let  $D_1$  and  $D_2$  be two close parallel copies of  $D$  in  $\text{Int } T$  and let  $B$  be a band which joins  $D_1$  to  $D_2$  as pictured in Figure 2.




FIGURE 2

Define  $C_1 = \partial D_1$ ,  $C_2 = \partial D_2$ ,  $D' = D_1 \cup B \cup D_2$  and  $C' = \partial D'$ . Choose  $\epsilon > 0$  small enough so that  $T \times [0, \epsilon]$  is contained in the interior of  $W_2$ . Then  $W_1$  is defined to be a regular neighborhood of

$$T \times [-1, 0] \cup C' \times [0, \epsilon] \cup B \times \{\epsilon\} \cup (C_1 \cup C_2) \times [\epsilon, 1] \cup (D_1 \cup D_2) \times \{1\}.$$

The regular neighborhood should be chosen so that  $W_1 \subset \text{Int } W_2$ . Notice that  $D'$  is isotopic to  $D$  in  $T$ , so  $W_1 \cong W_2$ . In fact,  $W_1$  is embedded nicely enough so that there is an isotopy  $h_t : S^4 \rightarrow S^4$ ,  $0 \leq t \leq 1$ , with  $h_0 = \text{id}$ , and  $h_1(W_1) = W_2$ . We recursively define  $W_n$ ,  $n \geq 3$ , by  $W_n = h_1(W_{n-1})$  and define  $W = \bigcup_{n=1}^{\infty} W_n$ .

It is clear that  $W_n \searrow S^1 \vee S^2$  and that the inclusion  $W_n \subset W_{n+1}$  induces the identity map on the  $S^1$  factor. It follows that  $\pi_1(W) \cong \mathbb{Z}$ . The band  $B$  has a half-twist in it (see Figure 2) and so the inclusion induced map  $H_2(W_n) \rightarrow H_2(W_{n+1})$  is trivial. Thus  $H_2(W) = 0$ . However, we will see that the fact that the band goes around the  $S^1$ -factor of  $W_{n+1}$  makes  $\pi_2(W)$  nontrivial.

In order to describe  $\pi_2(W)$  more explicitly, we fix some notation. We use  $J$  to denote  $\pi_1(W)$ ; so  $J \cong \mathbb{Z}$  and we write a typical element of  $J$  as  $t^n$ , where  $t$  is a generator of  $J$  and  $n \in \mathbb{Z}$ . Let  $\Lambda$  denote the group ring  $\mathbb{Z}[J]$ . Then  $\Lambda$  consists of all Laurent polynomials in  $t$  with integer coefficients. Since  $W_n \searrow S^1 \vee S^2$ ,  $\pi_2(W_n)$  is naturally isomorphic to  $\Lambda$ . The important thing to notice is that it is possible to choose a  $\Lambda$ -generator for  $\pi_2(W_n)$  in such a way that the inclusion map  $W_n \hookrightarrow W_{n+1}$  induces multiplication by  $t - 1$ ; i.e., the

diagram

$$\begin{array}{ccc}
 \pi_2(W_n) & \longrightarrow & \pi_2(W_{n+1}) \\
 \cong \downarrow & & \downarrow \cong \\
 \Lambda & \xrightarrow{t-1} & \Lambda
 \end{array}$$

commutes. It is easiest to see this by looking at the universal covers  $\widetilde{W}_n \subset \widetilde{W}_{n+1}$ .

It is now clear that  $\pi_2(W) \neq 0$ . In particular, the generator of  $\pi_2(W_1)$  represents  $(t-1)^n$  in  $\pi_2(W_{n+1})$ . Thus the generator of  $\pi_2(W_1)$  does not die in  $\pi_2(W_n)$  for any  $n$  and therefore does not die in  $\pi_2(W)$ .

We now turn our attention to the complements. Define  $Q_n = \overline{S^4 - W_n}$  and  $X = \bigcap_{n=1}^\infty Q_n = S^4 - W$ . The first thing to notice is that, since  $W_n$  is a regular neighborhood of  $S^1 \vee S^2$  and both factors are unknotted in  $S^4$ , we have that  $Q_n \searrow S^2 \vee S^1$ ; the  $S^2$ -factor of  $Q_n$  links the  $S^1$ -factor of  $W_n$  and the  $S^2$ -factor of  $Q_n$  links the  $S^1$ -factor of  $W_n$ . The  $S^1$ -factor of  $Q_2$  can be pictured as a loop  $A_2$  in the level  $\mathbb{R}^3 \times \{\frac{1}{2}\}$  such that  $A_2$  links  $C \times \{\frac{1}{2}\}$  and the  $S^2$ -factor can be pictured as the suspension in  $\mathbb{R}^4$  of a circle  $B_2$  in  $\mathbb{R}^3 \times \{\frac{1}{2}\}$  such that  $B_2$  links  $T \times \{\frac{1}{2}\}$ . This is indicated schematically in Figure 3.




FIGURE 3

In general,  $Q_n$  is a regular neighborhood of  $A_n \cup \mathcal{S}(B_n)$ , where  $A_n \cup B_n$  is a figure-8 in a level,  $\mathcal{S}$  denotes suspension, and all the  $B_n$ 's are the same. We wish to understand the embedding of  $Q_{n+1}$  in  $Q_n$ . We concentrate on  $Q_2 \subset Q_1$ . In Figure 4 we show  $A_1, A_2$  and  $B_1 = B_2$ .




FIGURE 4

Now  $A_1$  is the spine of the complement of  $C'$ . Thus, to understand the embedding of  $Q_2$  in  $Q_1$ , we must redraw the picture so that  $C'$  and  $A_1$  are straightened out. This is shown in Figure 5.




FIGURE 5

We are now in a position to give a geometric description of  $X = S^4 - W$ . We see that  $X = \bigcap_{n=1}^{\infty} Q_n$  where each  $Q_n$  a regular neighborhood of a copy of  $S^1 \vee S^2$ . Each  $S^1 \vee S^2 = A_n \vee \mathcal{S}(B_n)$  where  $A_n \cup B_n$  is a figure-8 in a hyperplane. Furthermore,  $B_{n+1} = B_n$ . The embedding of  $A_{n+1} \cup B_{n+1}$  in a neighborhood of  $A_n \cup B_n$  is indicated in Figure 6.




FIGURE 6

We now re-embed the  $Q_n$ 's so that they are closer and closer regular neighborhoods of the spines  $A_n \vee \mathcal{S}(B_n)$ . After that adjustment,  $X$  consists of a locally flat 2-sphere,  $\mathcal{S}(B_1)$ , together with a 1-dimensional continuum which we call Wh; i.e.,  $X = \mathcal{S}(B_1) \cup \text{Wh}$  and  $\mathcal{S}(B_1) \cap \text{Wh}$  consists of a single point. Since  $A_{n+1}$  is null-homotopic in a regular neighborhood of  $A_n$ , Wh is cell-like. In fact, Wh is cellular in  $S^4$  since it is the intersection of a sequence of neighborhoods with 1-dimensional spines. We can therefore shrink Wh to a point and the resulting manifold is still homeomorphic to  $S^4$ . We define  $\Sigma_1$  to be the image of  $X$  in the decomposition space  $S^4/\text{Wh}$ . Since  $S^4/\text{Wh} - \Sigma_1 = S^4 - X = W$ , it is clear that  $\Sigma_1$  has the properties required of Example 1.1.  $\square$

*Remark 1.* It should be noted that  $\Sigma_1$  is locally flat except at the image of Wh; at that point  $\Sigma_1$  cannot be locally flat. In fact,  $\Sigma_1$  cannot even be locally 1-*alg* at the exceptional point. This is because Hollingsworth and Rushing [9] have shown that any 2-sphere which is locally 1-*alg* at every point is globally 1-*alg*. Thus Corollary 3.5 would be contradicted if  $\Sigma_1$  were locally 1-*alg* at the exceptional point.  $\square$

*Remark 2.* Our construction contains a counterexample to the proof of [6, Lemma 9.3B]. The manifolds  $\{W_n\}$  above satisfy conditions (1)–(3) on p. 142 of [5] as well as (i)–(iv)

on p. 143, but it is not the case that the inclusion induced map  $\pi_2(W_n) \rightarrow \pi_2(W_{n+k})$  is trivial for any  $k \geq 1$ . The error in [6] is the assertion (on p. 146) that “condition (iv) is equivalent to ...  $I(\text{image}) = \text{image}$ .” In general it will only be the case that, in the terminology of [6],  $\text{image} \subset I(\text{whole group})$ . This is analyzed more carefully in the proof of Theorem 3.1, below.  $\square$

*Remark 3.* There exists a high dimensional version of our example. Let  $X = \mathcal{S}^{n-3}(B_1) \cup \text{Wh} \subset S^n$ , where  $B_1$  and Wh are the subsets of  $S^3$  described above and  $\mathcal{S}^{n-3}(B_1)$  denotes the  $(n-3)$ -fold suspension of  $B_1$ . Then

$$\Sigma^{n-2} = X/\text{Wh} \subset S^n/\text{Wh} \cong S^n$$

is a topologically embedded  $(n-2)$ -sphere in  $S^n$ ,  $n \geq 4$ . The complement of  $\Sigma^{n-2}$  has the property that  $\pi_1(S^n - \Sigma^{n-2}) \cong \mathbb{Z}$  and  $\pi_i(S^n - \Sigma^{n-2}) = 0$  for  $2 \leq i \leq n-3$ , but  $\pi_{n-2}(S^n - \Sigma^{n-2})$  is nontrivial. This contrasts sharply with Levine’s Theorem [10] which asserts the following: If  $S^{n-2}$  is a smooth  $(n-2)$ -sphere in  $S^n$  such that  $\pi_1(S^n - S^{n-2}) \cong \mathbb{Z}$  and  $\pi_i(S^n - S^{n-2}) = 0$  for  $2 \leq i < n/2$ , then  $S^{n-2}$  is unknotted,  $n \geq 6$ .  $\square$

*Remark 4.* The embedding  $h_0 : \mathbb{R}^2 \rightarrow \mathbb{R}^4$  of Example 1.2 is constructed by removing the point Wh from  $(S^4, \Sigma_1)$ . More generally, we can remove the point Wh from  $\Sigma^{n-2} \subset S^n$  (described in Remark 3, above), to construct an interesting example of an embedding of  $\mathbb{R}^{n-2}$  in  $\mathbb{R}^n$ . Again, this example contrasts sharply with Levine’s Theorem.  $\square$

**Example 1.2’.** *There exists a proper, smooth embedding  $h : \mathbb{R}^{n-2} \rightarrow \mathbb{R}^n$ ,  $n \geq 4$ , such that  $\pi_1(\mathbb{R}^n - h(\mathbb{R}^{n-2})) \cong \mathbb{Z}$  and  $\pi_i(\mathbb{R}^n - h(\mathbb{R}^{n-2})) = 0$  for  $2 \leq i \leq n-3$  but  $\pi_{n-2}(\mathbb{R}^n - h(\mathbb{R}^{n-2})) \neq 0$ .*

## 2. THE SECOND EXAMPLE

In this section we construct Example 2.1. The method of constructing the local example from the global one is the same as that described by Daverman on p. 372 of [3].

Let  $D'$  be a 4-cell in  $S^4$  such that  $D'$  intersects  $\Sigma_1$  in a locally flat disk missing the exceptional point of  $\Sigma_1$  and let  $D = \overline{S^4 - D'}$ . Then  $(D, D \cap \Sigma_1)$  is a knotted topological  $(4,2)$ -ball pair. Remove from  $S^4$  a sequence  $D_1, D_2, D_3, \dots$  of pairwise disjoint 4-cells such that, for each  $n$ ,  $(D_n, D_n \cap S^2)$  is an unknotted  $(4,2)$ -ball pair and  $\{D_n\}_{n=1}^\infty$  converges to a point  $x \in S^2$ . Replace each  $(D_n, D_n \cap S^2)$  with a  $(4,2)$ -ball pair  $(D_n, C_n)$  such that  $\partial C_n = \partial(D_n \cap S^2)$  and  $(D_n, C_n) \cong (D, D \cap \Sigma_1)$ . Then let

$$\Sigma_2 = \left[ S^2 - \left( \bigcup_{n=1}^{\infty} (D_n \cap S^2) \right) \right] \cup \left( \bigcup_{n=1}^{\infty} C_n \right).$$

Since  $\pi_1(D - \Sigma_1) \cong \mathbb{Z}$ , Van Kampen's theorem implies that  $\Sigma_2$  is locally 1-*alg* at  $x$ . A Mayer-Vietoris argument in the universal cover of  $S^4 - \Sigma_2$  shows that the local  $\pi_2$  of  $S^4 - \Sigma_2$  is not trivial at  $x$ . Thus  $\Sigma_2$  has the properties required of Example 2.1.  $\square$

In the Introduction we described an example of a PL embedding of  $S^{n-2}$  into  $S^n$ ,  $n \geq 6$ , which shows that locally 1-*alg* does not imply locally homotopically unknotted. We have just constructed an example of such an embedding for  $n = 4$  as well. For the sake of completeness, we now construct an example for  $n = 5$ . Let  $\Sigma^3 \subset S^5$  be a locally flat PL 3-sphere such that  $\pi_1(S^5 - \Sigma^3) \cong \mathbb{Z}$  and  $\pi_2(S^5 - \Sigma^3) \neq 0$ . By removing an unknotted ball pair from  $(S^5, \Sigma^3)$ , we obtain a knotted ball pair  $(D^5, D^5 \cap \Sigma^3)$ . Proceeding as in the construction of Example 2.1, we start with the unknotted sphere pair  $(S^5, S^3)$ , remove a null-sequence of balls from it, and replace each with a copy of  $(D^5, D^5 \cap \Sigma^3)$  to produce a topological embedding of  $S^3$  into  $S^5$  having the following properties.

**Example 2.2.** *There exists a topological embedding  $h_3 : S^3 \rightarrow S^5$  and a point  $x_3 \in h_3(S^3)$  such that  $h_3(S^3)$  is locally 1-*alg* at  $x_3$  but  $h_3(S^3)$  is not locally homotopically unknotted at  $x_3$ .*

Notice that  $h_3(S^3)$  is locally flat and PL at every point other than  $x_3$  and that it is the local  $\pi_2$  that is bad at  $x_3$ . We could construct an example with bad local  $\pi_3$  by applying the infinite construction to the example mentioned in Remark 3 at the end of §1. The example constructed in that way would have an infinite sequence of non-locally flat points. Obviously our 5-dimensional examples are not PL as the high dimensional examples were. Our next theorem shows that there can be no PL embedding of  $S^3$  in  $S^5$  having the properties mentioned in Example 2.2, but that PL embeddings of  $S^3$  into  $S^5$  follow the low-dimensional pattern.

**Theorem 2.3.** *If  $M^3$  is a 3-dimensional PL submanifold of the PL 5-manifold  $W^5$  and  $M^3$  is locally 1-*alg* at  $x$  for every  $x \in M^3$ , then  $M^3$  is (topologically) locally flat.*

*Proof.* Let  $x \in M^3$  and let  $\sigma$  be the simplex of  $M^3$  such that  $x \in \text{Int } \sigma$ . If  $\text{lk } \sigma$  denotes the link of  $\sigma$  in  $W^5$ , then  $(\text{lk } \sigma, \text{lk } \sigma \cap M^3)$  is a PL sphere pair of dimension  $(4 - \dim \sigma, 2 - \dim \sigma)$ . The fact that  $M^3$  is locally 1-*alg* at  $x$  implies that  $\pi_1(\text{lk } \sigma - M^3) \cong \mathbb{Z}$ . As long as  $\dim \sigma \geq 2$ , it is obvious that  $(\text{lk } \sigma, \text{lk } \sigma \cap M^3)$  is an unknotted sphere pair. In case  $\dim \sigma = 1$ , this follows from the Loop Theorem [14] and in case  $\dim \sigma = 0$ ,  $(\text{lk } \sigma, \text{lk } \sigma \cap M^3)$  is a topologically unknotted (4,2)-sphere pair by [5, Theorem 6].  $\square$

### 3. ASPHERICAL 2-SPHERE COMPLEMENTS

We next turn our attention to the problem of proving a positive theorem regarding aspherical 2-spheres in  $S^4$ . For the remainder of this section, we let  $\Sigma$  denote a 2-sphere topologically embedded in  $S^4$  and  $W = S^4 - \Sigma$ . We assume that  $\Sigma$  is embedded in such a

way that  $\pi_1(W) \cong \mathbb{Z}$ . In §1 we showed that  $\pi_2(W)$  may be nontrivial and thus  $W$  is not necessarily aspherical. In this section we give conditions which imply that  $W$  is aspherical. It should be noted that, by the Whitehead theorem,  $W$  has the homotopy type of  $S^1$  if and only if  $\pi_i(W) = 0$  for  $i \geq 2$ . But the Hurewicz theorem implies that  $\pi_i(W) = 0$  for  $i \geq 2$  if  $H_i(\widetilde{W}) = 0$  for  $i \geq 2$ , where  $\widetilde{W}$  is the universal cover of  $W$ . Thus we will concentrate in this section on the problem of computing the homology of  $\widetilde{W}$ . The main result of this section (Theorem 3.1) shows that any nontrivial higher homology groups must appear at the  $H_2$  level and that this will only happen if  $H_2$  is quite complicated. Our other results in this section show that most of the algebraic properties of Example 1.1 are common to all examples.

We begin the proof of Theorem 3.1 by constructing two sequences of submanifolds of  $S^4$  which will be useful in the proofs below. Let  $U_1 \supset U_2 \supset U_3 \supset \dots$  be a sequence of compact connected PL manifold neighborhoods of  $\Sigma$  such that  $U_{n+1} \subset \text{Int} U_n$  for each  $n$  and  $\bigcap_{n=1}^{\infty} U_n = \Sigma$ . Since  $\Sigma$  is an ANR, there is a retraction  $r_n : U_n \rightarrow \Sigma$  for large  $n$ . We may assume that  $U_n$  is chosen in such a way that  $r_n$  is homotopic to the inclusion in  $U_{n-1}$ . Since  $\Sigma$  is 2-dimensional,  $\Sigma$  does not separate any neighborhood and so we may also assume that  $\partial U_n$  is connected for each  $n$ . (If not, connect up the boundary components with arcs lying in  $U_n - \Sigma$  and remove regular neighborhoods of the arcs.) Define  $W_n = \overline{S^4 - U_n}$ . It follows from Alexander duality that  $H_2(W) \cong H^1(\Sigma) = 0$ . Each  $H_2(W_n)$  is finitely generated (since  $W_n$  is a compact PL manifold) so we may also assume that the inclusion induced map of  $H_2(W_n)$  into  $H_2(W_{n+1})$  is trivial. In addition, we use the fact that  $\pi_1(W) \cong \mathbb{Z}$  to arrange that any loop in  $W_n$  which is null-homotopic in  $W$  is null-homotopic in  $W_{n+1}$  and that  $\pi_1(W_n) \rightarrow \pi_1(W)$  is onto. Let  $p : \widetilde{W} \rightarrow W$  denote the covering projection.

**Lemma 3.6.** *For each  $n$ , the image of  $H_1(U_{n+1} - \Sigma)$  in  $H_1(U_n - \Sigma)$  is canonically isomorphic to  $H_1(W) \cong \mathbb{Z}$ . It follows that both  $p^{-1}(U_n - \Sigma)$  and  $p^{-1}(\partial U_n)$  are connected and that  $\widetilde{W}$  has one end.*

*Proof.* Fix  $n$ . Let  $\gamma : H_1(U_{n+1} - \Sigma) \rightarrow H_1(U_n - \Sigma)$  and  $\eta : H_1(U_n - \Sigma) \rightarrow H_1(W)$  denote the inclusion induced homomorphisms. We will show that  $\eta|_{\text{im } \gamma} : \text{im } \gamma \rightarrow H_1(W) \cong \mathbb{Z}$  is an isomorphism. We begin by showing that  $\text{im } \gamma$  is cyclic. Consider the commutative diagram

$$\begin{array}{ccccc} H_1(U_{n+1} - \Sigma) & \longrightarrow & H_1(U_{n+1}) & & \\ & & \gamma \downarrow & & \downarrow \delta \\ H_2(U_n, U_n - \Sigma) & \xrightarrow{\alpha} & H_1(U_n - \Sigma) & \xrightarrow{\beta} & H_1(U_n). \end{array}$$

Since the inclusion  $U_{n+1} \hookrightarrow U_n$  factors, up to homotopy, through a retraction onto  $\Sigma$ , we have that  $\delta = 0$ . Therefore  $\beta \circ \gamma = 0$  and so  $\text{im } \gamma \subset \ker \beta = \text{im } \alpha$ . But  $H_2(U_n, U_n - \Sigma) \cong \mathbb{Z}$  by Alexander duality, so  $\text{im } \gamma$  is cyclic.

Now consider the commutative diagram

$$\begin{array}{ccccc} H_1(U_{n+1} - \Sigma) & \xrightarrow{\zeta} & H_1(S^4 - \Sigma) & \longrightarrow & H_1(S^4 - \Sigma, U_{n+1} - \Sigma) \\ \gamma \downarrow & & \parallel & & \\ H_1(U_n - \Sigma) & \xrightarrow{\eta} & H_1(S^4 - \Sigma) & & \end{array}$$

By excision,  $H_1(S^4 - \Sigma, U_{n+1} - \Sigma) \cong H_1(S^4, U_{n+1})$ . Since  $U_{n+1}$  is connected, it follows that  $H_1(S^4, U_{n+1}) = 0$ . Therefore  $\zeta$  is onto and hence  $\eta \circ \gamma$  is onto. This implies that  $\eta|_{\text{im } \gamma} : \text{im } \gamma \rightarrow H_1(S^4 - \Sigma) \cong \mathbb{Z}$  is an isomorphism since the only onto homomorphisms from a cyclic group to  $\mathbb{Z}$  are isomorphisms.

In order to see that each  $p^{-1}(U_n - \Sigma)$  is connected, it is enough to note that the composition  $\pi_1(U_n - \Sigma) \rightarrow H_1(U_n - \Sigma) \rightarrow H_1(W) \cong \pi_1(W)$  is onto. A similar proof shows that  $p^{-1}(\partial U_n)$  is connected. Since  $\partial U_n$  is connected, the only other ingredient we need is a representative of the generator  $t$  of  $H_1(W)$  in  $\partial U_n$ . We have already seen that there is a representative of  $t$  in  $U_n - \Sigma$  and the choice of  $W_n$  shows that there is a representative of  $t$  in  $W_n$ . Hence, a Mayer-Vietoris sequence argument shows that there is a class in  $H_1(\partial U_n)$  which is homologous in  $W$  to  $t$ .

For each  $n$ , we can find a compact connected polyhedron  $K_n \subset \widetilde{W}$  such that  $W_n \subset p(K_n)$ . A typical neighborhood of  $\infty$  in  $\widetilde{W}$  has the form

$$V_{n,k} = p^{-1}(U_n - \Sigma) \cup \left( \bigcup_{|j| \geq k} t^j K_n \right)$$

Since  $V_{n,k}$  is connected for every  $n$  and  $k$ ,  $\widetilde{W}$  has one end.  $\square$

As in §1, we use  $J$  to denote  $\pi_1(W) \cong \mathbb{Z}$ ,  $\Lambda$  to denote the integral group ring  $\mathbb{Z}[J]$ , and  $t$  to denote a generator of  $J$ . Let  $p : \widetilde{W} \rightarrow W$  be the universal cover. If  $A$  is a subset of  $W$  such that  $\pi_1(A) \rightarrow \pi_1(W) \cong \mathbb{Z}$  is onto, we write  $H_k(A; \Lambda)$  for  $H_k(p^{-1}(A); \mathbb{Z})$ . Here we are thinking of  $\Lambda$ -coefficients as local coefficients in the sense of Steenrod [15]. In case  $A$  is a compact  $n$ -dimensional submanifold of  $W$ , there is a Poincaré duality theorem for  $H_*(A; \Lambda)$ . It asserts that  $H_k(A; \Lambda) \cong H^{n-k}(A, \partial A; \Lambda)$ . In order to avoid confusion it should be remembered that  $H^i(A; \Lambda)$  is the same as  $H_c^i(p^{-1}(A); \mathbb{Z})$ .

**Lemma 3.7.** *For each  $n$ ,  $H_3(W_n; \Lambda) = 0$ ,  $H^3(W_n; \Lambda) = 0$ , and  $H_2(W_n; \Lambda)$  is a submodule of a finitely generated free  $\Lambda$ -module.*

*Remark.* Since  $H_*(\widetilde{W})$  is the direct limit of the images of the groups  $H_*(W_n; \Lambda)$ , we can conclude from the Lemma that every complement  $W$  of the sort studied in this section will possess many of the same algebraic properties that the complement of Example 1.1 had. In particular,  $H_i(\widetilde{W}) = 0$  for  $i \geq 3$  and  $H_2(\widetilde{W})$  is the direct limit of a sequence of

modules, each of which is a submodule of a finitely generated free module. In Theorem 3.1 we show that the only way in which  $H_2(\widetilde{W})$  itself can be a submodule of a free module is if it is trivial.  $\square$

*Proof of Lemma 3.7.* First observe that the pair  $(W, U_n - \Sigma)$  is 1-connected. By Poincaré duality and excision we have  $H^3(W_n; \Lambda) \cong H_1(W_n, \partial W_n; \Lambda) \cong H_1(W, U_n - \Sigma; \Lambda) = 0$ . Similarly,  $H_3(W_n; \Lambda) \cong H^1(W_n, \partial W_n; \Lambda) \cong H^1(W, U_n - \Sigma; \Lambda) = 0$ .

Since  $W_n$  collapses to a compact 3-dimensional polyhedron, the chain complex for  $H_*(W_n; \Lambda)$  can have the form

$$0 \rightarrow C_3 \xrightarrow{\partial_3} C_2 \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0$$

Because  $H_3(W_n; \Lambda) = 0$ ,  $\partial_3$  is injective and  $\text{im } \partial_3$  is a finitely generated free  $\Lambda$ -module. We can think of  $[\partial_3]$  as an element of  $H^3(W_n; \text{im } \partial_3)$ . Moreover  $H^3(W_n; \text{im } \partial_3) = 0$  since  $\text{im } \partial_3$  is a finitely generated free  $\Lambda$ -module and  $H^3(W_n; \Lambda) = 0$ . So there exists  $\theta \in \text{Hom}_\Lambda(C_2, \text{im } \partial_3)$  such that  $\partial_3 = \delta_3(\theta) = \theta \circ \partial_3$ . Let  $Z_2 = \ker \partial_2$ . Then  $\theta|_{Z_2}$  splits the sequence

$$0 \rightarrow \text{im } \partial_3 \hookrightarrow Z_2 \rightarrow H_2(W_n; \Lambda) \rightarrow 0.$$

Thus  $H_2(W_n; \Lambda)$  is isomorphic to a submodule of  $Z_2 \subset C_2$ .  $\square$

*Proof of Theorem 3.1.* It is obvious that condition (1) implies each of the others and that condition (2) implies each of (3) and (4). It follows from Lemma 3.7 along with the Whitehead and Hurewicz theorems that (2) implies (1). We will show that (3) implies (2) and that (4) implies (2).

Suppose, first, that  $H_2(\widetilde{W})$  is a submodule of a free  $\Lambda$ -module. The short exact sequence

$$0 \rightarrow C_*(\widetilde{W}) \xrightarrow{t-1} C_*(\widetilde{W}) \xrightarrow{p_*} C_*(W) \rightarrow 0$$

of chain complexes gives rise to the exact sequence

$$0 = H_3(W) \rightarrow H_2(\widetilde{W}) \xrightarrow{t-1} H_2(\widetilde{W}) \xrightarrow{p_*} H_2(W) = 0$$

of homology groups. Thus  $(t-1) : H_2(\widetilde{W}) \rightarrow H_2(\widetilde{W})$  is onto. This means that each element of  $H_2(\widetilde{W})$  is infinitely divisible by  $(t-1)$ , but the only element of a free module with that property is 0. Thus  $H_2(\widetilde{W})$  cannot be a submodule of a free module unless it is trivial.

Second, suppose that  $H_2(\widetilde{W})$  is finitely generated over  $\Lambda$ . We fix a finite set of generators and choose  $W_1$  in such a way that  $p^{-1}(W_1)$  contains representatives of this set of generators for  $H_2(W; \Lambda)$ . Each  $H_2(W_n; \Lambda)$  is finitely generated, so we can choose  $W_{n+1}$  large enough that each element of  $H_2(W_n; \Lambda)$  is homologous in  $W_{n+1}$  to some combination of the fixed

set of generators. This will mean that the image of  $H_2(W_n; \Lambda)$  in  $H_2(W_{n+1}; \Lambda)$  will be equal to the image of  $H_2(W_1; \Lambda)$  in  $H_2(W_{n+1}; \Lambda)$ . In this way we arrange that, for each  $n$ , the image of  $H_2(p^{-1}(W_n))$  in  $H_2(p^{-1}(W_{n+2}))$  is the same as that of  $H_2(p^{-1}(W_{n+1}))$  in  $H_2(p^{-1}(W_{n+2}))$ . As in the previous paragraph, the rows in the following diagram are exact.

$$\begin{array}{ccccc}
 H_2(p^{-1}(W_n)) & \xrightarrow{t-1} & H_2(p^{-1}(W_n)) & \xrightarrow{p_*} & H_2(W_n) \\
 \downarrow & & \alpha_n \downarrow & & \downarrow 0 \\
 H_2(p^{-1}(W_{n+1})) & \xrightarrow{t-1} & H_2(p^{-1}(W_{n+1})) & \xrightarrow{p_*} & H_2(W_{n+1}) \\
 \downarrow & & \alpha_{n+1} \downarrow & & \downarrow 0 \\
 H_2(p^{-1}(W_{n+2})) & \xrightarrow{t-1} & H_2(p^{-1}(W_{n+2})) & \xrightarrow{p_*} & H_2(W_{n+2})
 \end{array}$$

This means that  $\text{im } \alpha_n \subset (t-1)(H_2(p^{-1}(W_{n+1})))$  and so

$$\text{im } (\alpha_{n+1} \circ \alpha_n) \subset (t-1)\text{im } \alpha_{n+1} = (t-1)\text{im } (\alpha_{n+1} \circ \alpha_n)$$

Therefore every element of  $\text{im } (\alpha_{n+1} \circ \alpha_n)$  is infinitely divisible by  $(t-1)$ . By Lemma 3.7,  $H_2(p^{-1}(W_{n+2}))$  is a submodule of a free  $\Lambda$ -module and thus  $\text{im } (\alpha_{n+1} \circ \alpha_n)$  is a submodule of a free  $\Lambda$ -module. This means that  $\text{im } (\alpha_{n+1} \circ \alpha_n) = 0$  and hence  $H_2(\widetilde{W}) = 0$ . The proof of the theorem is complete.  $\square$

*Proof of Corollary 3.4.* Since  $\pi_1(\epsilon) = \mathbb{Z}$  and the natural map  $\pi_1(\epsilon) \rightarrow \pi_1(W)$  is an isomorphism (Lemma 3.6),  $\widetilde{W}$  has one simply connected end. This means that  $H^3(\widetilde{W}) \cong H_1^\infty(\widetilde{W}) = 0$ . ( $H_*^\infty(\widetilde{W})$  denotes the homotopy based on infinite chains.) It follows that  $H_2(\widetilde{W})$  is a projective module over  $\Lambda$  (by the argument of [11, Lemma 2.4] or [18, Lemma 2.1]) and so the Theorem applies.  $\square$

#### 4. SURFACES WHICH ARE LOCALLY 1-ALG AT EVERY POINT

In this final section of the paper we consider the question of whether or not a topologically embedded surface in a 4-manifold which is locally 1-alg at every one of its points is locally homotopically unknotted (and therefore locally flat). We are not able to answer that question in general, but do prove a positive result in the important special case in which the surface is a 2-sphere. In this section we will make use of the following technical lemma from homological algebra.

**Lemma 4.5.** *Suppose  $W$  is a PL manifold with  $\pi_1(W) \cong \mathbb{Z}$  and  $A \subset B$  are two polyhedral subsets of  $W$  such that the pairs  $(W, W - A)$  and  $(W, W - B)$  are  $(q-1)$ -connected for some  $q$ . If the inclusion induced homomorphism  $H_q(W, W - B; \Lambda) \rightarrow H_q(W, W - A; \Lambda)$  is the*

zero homomorphism, then the restriction homomorphism  $H^q(W, W - A; \Lambda) \rightarrow H^q(W, W - B; \Lambda)$  is also zero.

*Sketch of proof.* First notice that  $(W, W - A)$  and  $(W, W - B)$  are simple homotopy equivalent to pairs with no relative  $n$ -cells,  $n < q$  (cf. [18] Lemma 1.1). Thus the cellular chain complexes have the form

$$\begin{array}{ccccccc} \cdots & \longrightarrow & C_{q+1}(W, W - B; \Lambda) & \xrightarrow{\partial} & C_q(W, W - B; \Lambda) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & C_{q+1}(W, W - A; \Lambda) & \xrightarrow{\partial} & C_q(W, W - A; \Lambda) & \longrightarrow & 0 \end{array}$$

Take  $\text{Hom}_\Lambda(\cdot, \Lambda)$  of the diagram above. Then the proof is completed by a straightforward argument using the definitions of homology and cohomology.  $\square$

To begin the proof of Theorem 4.1, we assume that  $\Sigma$  is a 2-sphere topologically embedded in the interior of the 4-manifold  $M^4$  in such a way that the self intersection number is 0. Our next two lemmas do not require the hypothesis that  $\Sigma$  be locally 1-*alg* at every point, but only the weaker hypothesis that  $\Sigma$  is globally 1-*alg*. Thus that is all we will assume for the moment. (It is shown in [9] that if  $\Sigma$  is locally 1-*alg* at every point, then  $\Sigma$  is globally 1-*alg*. Hence the global hypothesis is strictly weaker than the local one.)

Since  $\Sigma$  is globally 1-*alg* and the self intersection number is 0, the main theorem of [12] implies that there is a collar  $C$  of the end of  $M - \Sigma$ . Specifically, there exists a compact neighborhood  $N$  of  $\Sigma$  and a homeomorphism  $\phi : S^1 \times S^2 \times [0, 1) \rightarrow C = N - \Sigma$ . Let  $p : \tilde{C} \rightarrow C$  be the universal cover of  $C$ . As before, we use  $J$  to denote  $\pi_1(C)$  and  $\Lambda = \mathbb{Z}[J]$ . For  $0 < a < 1$  we use  $C_a$  to denote the subcollar  $C_a = \phi(S^1 \times S^2 \times [a, 1))$ .

**Lemma 4.6.** *Suppose  $B$  is a PL 4-ball in the interior of  $C \cup \Sigma$ ,  $0 < a < 1$ , and  $B - (\text{Int } C_a \cup \Sigma)$  is PL with respect to the PL structure on  $C$  induced by  $\phi$ . If  $P = \overline{B - (C_a \cup \Sigma)}$ , then  $H_3(P; \Lambda) = 0$ ,  $H^3(P; \Lambda) = 0$ , and  $H_2(P; \Lambda)$  is a submodule of a finitely generated free  $\Lambda$ -module.*

*Proof.* We first prove that  $C - P$  is connected. Notice that  $C - P = C_a \cup (C - B)$  and that it is obvious that  $C_a$  is connected. Given two points  $y$  and  $z$  in  $C - B$ , there exists an arc  $A \subset C$  joining them. Put  $A$  in general position with respect to  $\partial B$ . Then  $A \cap \partial B$  will consist of a finite number of points which divide  $A$  into a finite number of subarcs, each of which is entirely inside  $B$  or entirely outside  $B$ . We replace each subarc of  $A$  which lies inside  $B$  with an arc lying on  $\partial B$ . The new arc from  $y$  to  $z$  can be pushed into the complement of  $B$ . Since  $\Sigma$  is 2-dimensional,  $\Sigma$  does not separate any open set so we can adjust the arc slightly to miss  $\Sigma$  as well. Thus  $C - B$  is connected and so is  $C - P$ .

Since  $\pi_1(C - P) \rightarrow \pi_1(C)$  is onto, we conclude that the pair  $(C, C - P)$  is 1-connected. The Lemma can now be proved by the same argument as was used to prove Lemma 3.7.  $\square$

**Lemma 4.7.** *Fix  $x \in \Sigma$ . Suppose  $B_1$  is a 4-ball in  $C \cup \Sigma$  such that  $x \in \text{Int } B_1$ . Let  $D$  be a disk on  $\Sigma$  such that  $x \in \text{Int } D$  and  $D \subset \text{Int } B_1$  and let  $B_2$  be a 4-ball such that  $x \in \text{Int } B_2$ ,  $B_2 \cap \Sigma \subset \text{Int } D$ , and  $B_2 \subset \text{Int } B_1$ . For every  $a$ ,  $0 < a < 1$ , there exists a number  $b$ ,  $a < b < 1$ , such that if  $P_1 = \overline{B_1 - (C_b \cup \Sigma)}$  and  $P_2 = \overline{B_2 - (C_a \cup \Sigma)}$ , then the inclusion induced homomorphism  $H_1(C - \text{Int } P_1; \Lambda) \rightarrow H_1(C - \text{Int } P_2; \Lambda)$  is trivial.*

*Proof.* Let  $x, B_1, D, B_2$ , and  $a$  be given. Choose  $V$  to be a connected PL manifold neighborhood of  $\partial D$  in  $\text{Int}((B_1 - B_2) \cap C_a)$ . Then choose  $b > a$  so that  $V$  separates  $C_b \cap B_2$  from  $C_b \cap (C - B_1)$  in  $C_b$ . Define  $P_1 = \overline{B_1 - (C_b \cup \Sigma)}$  and  $P_2 = \overline{B_2 - (C_a \cup \Sigma)}$ . We must show that each loop in  $p^{-1}(C - \text{Int } P_1)$  is null-homologous in  $p^{-1}(C - \text{Int } P_2)$ .

Let  $\gamma$  be a loop in  $p^{-1}(C - \text{Int } P_1)$  and let  $\gamma' = p(\gamma)$ . Put  $\gamma'$  in general position with respect to  $\partial B_1$ . Then  $\gamma' \cap \partial B_1$  will consist of a finite number of points. These points can be paired off so that each pair either bounds a subarc of  $\gamma'$  which lies outside  $B_1$  or a subarc which lies in  $C_b$ . Let  $y_1, y_2$  be a pair which bounds a subarc in  $C_b$ . We claim that there is an arc from  $y_1$  to  $y_2$  which lies in  $C_a - B_2$ . If the subarc of  $\gamma'$  which joins  $y_1$  to  $y_2$  lies outside  $B_2$ , we can use it. Otherwise the subarc must pass through  $V$  and so we can replace the portion from the first point of intersection with  $V$  to the last point of intersection with  $V$  with an arc in  $V - \Sigma$  to obtain the arc we need to prove the claim. Use these arcs to write  $\gamma' = \gamma_1 + \gamma_2$ , where  $\gamma_1 \subset C - B_2$  and  $\gamma_2 \subset C_a$ . By adding a multiple of the generator of  $\pi_1(C_a)$  to  $\gamma_1$  and its inverse to  $\gamma_2$ , we can arrange that both  $\gamma_1$  and  $\gamma_2$  are null-homotopic in  $C$ . This makes  $\gamma_2$  null-homotopic in  $C_a$ , so we have  $\gamma'$  homotopic to  $\gamma_1$  in  $C - \text{Int } P_2$ .

Now  $\gamma_1$  is null-homotopic in  $C$ . Since  $\partial B_2$  is simply connected, this null-homotopy can be cut off on  $\partial B_2$ . In other words, there exists a continuous function  $F : \Delta^2 \rightarrow (C - \text{Int } B_2) \cup \partial B_2$  such that  $F(\partial \Delta^2) = \gamma_1$ . There exist a finite number of pairwise disjoint disks  $\Delta_1, \dots, \Delta_n$  in  $\Delta^2$  such that

$$F^{-1}(\Sigma) \subset \bigcup_{i=1}^n \text{Int } \Delta_i$$

and  $F(\partial \Delta_i) \subset \text{Int } C_b$  for each  $i$ . Choose a basepoint  $z_0 \in \partial \Delta^2$  such that  $F(z_0) \in C_b$  and let  $A_1, \dots, A_n$  be a collection of oriented arcs in  $\Delta^2$  such that  $A_i$  is an arc from  $z_0$  to  $\partial \Delta_i$  and  $A_i \cap A_j = z_0$  for  $i \neq j$ . Let  $\delta_i$  denote the loop  $F(\partial \Delta_i)$  and let  $\alpha_i = F(A_i)$ . Then  $\gamma_1$  is homotopic in  $C - \text{Int } P_2$  to the loop  $\alpha_1 \delta_1 \alpha_1^{-1} \cdots \alpha_n \delta_n \alpha_n^{-1}$ . Let  $\beta_i$  be an arc in  $C_b$  which joins the ends of  $\alpha_i$ . By adding a multiple of the generator of  $\pi_1(C_b)$  to  $\beta_i$ , we can arrange that the loop  $\alpha_i \beta_i^{-1}$  is null-homotopic in  $C$ . Now  $\gamma_1$  is homologous in  $C - \text{Int } P_2$  to the loop  $\gamma_3 = \beta_1 \delta_1 \beta_1^{-1} \cdots \beta_n \delta_n \beta_n^{-1}$ . Furthermore, the fact that  $\alpha_i \beta_i^{-1}$  is null-homotopic in  $C$  means that this homology can be lifted to  $\tilde{C}$ . So  $\gamma$  is homologous in  $p^{-1}(C - \text{Int } P_2)$  to a lift  $\tilde{\gamma}_3$  of  $\gamma_3$ . This completes the proof because  $\tilde{\gamma}_3$  is contained in the simply connected space  $p^{-1}(C_b) \subset p^{-1}(C - P_2)$ .  $\square$

*Proof of Theorem 4.1.* Since  $\Sigma$  is locally 1-*alg* at each of its points, it is globally 1-*alg* [9]. Thus we may apply the two lemmas above. Fix  $x \in \Sigma$ . We wish to show that  $C$  is locally  $i$ -connected at  $x$  for every  $i \geq 2$ .

Let  $P_1$  and  $P_2$  be as in Lemma 4.7. Consider the diagram

$$\begin{array}{ccccc} H_2(C, C - \text{Int } P_1; \Lambda) & \longrightarrow & H_1(C - \text{Int } P_1; \Lambda) & & \\ & & \downarrow & & \downarrow \beta \\ H_2(C; \Lambda) & \xrightarrow{\alpha} & H_2(C, C - \text{Int } P_2; \Lambda) & \longrightarrow & H_1(C - \text{Int } P_2; \Lambda) \end{array}$$

Since  $\alpha$  is trivial and  $\beta$  is trivial (by Lemma 4.7), we have that the inclusion induced homomorphism  $H_2(C, C - \text{Int } P_1; \Lambda) \rightarrow H_2(C, C - \text{Int } P_2; \Lambda)$  is trivial. Moreover, since the pairs  $(C, C - \text{Int } P_1)$  and  $(C, C - \text{Int } P_2)$  are 1-connected, it follows from Lemma 4.5 that the restriction homomorphism  $H^2(C, C - \text{Int } P_2; \Lambda) \rightarrow H^2(C, C - \text{Int } P_1; \Lambda)$  is trivial. Therefore its dual, the inclusion induced homomorphism  $H_2(P_2; \Lambda) \rightarrow H_2(P_1; \Lambda)$ , is trivial as well.

Now let  $P_1 \supset P_2 \supset P_3 \supset \dots$  be a sequence of submanifolds of  $C$ , constructed as in Lemma 4.7. Then Lemmas 4.6 and 4.7 show that the inclusion induced homomorphisms  $H_i(P_k; \Lambda) \rightarrow H_i(P_{k-1}; \Lambda)$  are trivial for  $i \geq 2$ . It follows from the Eventual Hurewicz Theorem [4, Lemma 3.1] that there is an  $n$  such that  $\pi_i(P_k) \rightarrow \pi_i(P_{k-n})$  is trivial for  $i \geq 2$ .

Given a neighborhood  $U$  of  $x$ , we can find a 4-ball  $B_0$  such that  $x \in \text{Int } B_0$  and  $B_0 \subset U$ . Let  $B_0 \supset B_1 \supset \dots \supset B_n$  be 4-balls as in Lemma 4.7. Consider a map  $f : S^i \rightarrow B_n - \Sigma$ ,  $i \geq 2$ . Choose  $a$  so that  $f(S^i) \subset B_n - C_a$  and construct  $P_0, \dots, P_n$  from  $B_0, \dots, B_n$  as in Lemma 4.7 with all the subcollars involved being subcollars of  $C_a$ . Then  $f$  is null-homotopic in  $P_0$  by the previous paragraph. Hence  $f$  is null-homotopic in  $U - \Sigma$  and the proof is complete.  $\square$

*Proof of Corollary 4.3.* Pick a point  $x$  in  $N$ . There exists a disk  $D$  such that  $x \in \text{Int } D \subset D \subset N$  and  $N$  is locally flat in a neighborhood of  $\partial D$ . Let  $A$  be an annulus such that  $\partial D \subset A \subset D$  and  $N$  is locally flat at every point of  $A$ . Then there exists an embedding  $H : A \times B^2 \rightarrow M$  such that  $H(x, 0) = x$  for every  $x \in A$ . Choose a compact neighborhood  $W$  of  $D$  such that  $H(A \times B^2) \subset W, H(\partial D \times B^2) \subset \partial W$  and  $W \cap N = D$ . Form  $M'$  by attaching a 2-handle to  $W$  along  $\partial D \times B^2$  and let  $\Sigma \subset M'$  consist of  $D$  together with the core of the 2-handle. By changing the framing of the 2-handle, if necessary, we can arrange that the self-intersection number of  $\Sigma$  is 0. By Theorem 4.1,  $\Sigma$  is locally flat and thus  $N$  is locally flat at each point of  $\text{Int } D$ .  $\square$

*Proof of Corollary 4.4.* Remove a flat ball pair from a neighborhood of the locally flat point and then attach a 2-handle with appropriate framing so that the new 2-sphere has self-intersection number 0. Apply Theorem 4.1 to the new 2-sphere.  $\square$

## REFERENCES

1. J. Cannon, *ULC properties in neighborhood of embedded surfaces and curves in  $E^3$* , Can. J. Math. **25** (1973), 31–73.
2. T. A. Chapman, *Compact Hilbert cube manifolds and the invariance of Whitehead torsion*, Bull. Amer. Math. Soc. **79** (1973), 52–56.
3. R. J. Daverman, *Homotopy classification of complements of locally flat codimension two spheres*, Amer. J. Math. **98** (1976), 367–374.
4. S. C. Ferry, *Homotoping  $\epsilon$ -maps to homeomorphisms*, Amer. J. Math. **101** (1979), 567–582.
5. M. Freedman, *The disk theorem for four-dimensional manifolds*, Proceedings of the International Congress of Mathematicians, Polish Scientific Publishers, Warsaw, 1984, pp. 647–663.
6. M. H. Freedman and F. Quinn, *Topology of 4-manifolds*, Princeton University Press, Princeton, 1990.
7. C. Guillbault, *An open collar theorem for 4-manifolds*, Trans. Amer. Math. Soc. **331** (1992), 227–245.
8. P. S. Hirschhorn and J. G. Ratcliffe, *A simple proof of the algebraic unknotting of spheres in codimension two*, Amer. J. Math. **102** (1980), 489–491.
9. J. G. Hollingsworth and T. B. Rushing, *Homotopy characterization of weakly flat codimension 2 spheres*, Amer. J. Math. **98** (1976), 385–394.
10. J. Levine, *Unknotting spheres in codimension two*, Topology **4** (1965), 9–16.
11. V. T. Liem and G. A. Venema, *Characterization of knot complements in the 4-sphere*, Topology and its Appl. **42** (1991), 231–245.
12. ———, *Complements of 2-spheres in 4-manifolds*, Topology Hawaii (to appear).
13. J. Milnor, *Infinite cyclic coverings*, Conference on the Topology of Manifolds, Prindle, Weber and Schmidt, Boston, 1968, pp. 115–133.
14. C. D. Papakyriakopoulos, *On Dehn's lemma and the asphericity of knots*, Ann. Math. **66** (1957), 1–26.
15. N. E. Steenrod, *Homology with local coefficients*, Ann. of Math. **44** (1943), 610–627.
16. D. W. Sumners, *Homotopy torsion in codimension two knots*, Proc. Amer. Math. Soc. **24** (1970), 229–240.
17. ———, *On asphericity of knots of  $S^2$  in  $S^4$* , (unpublished manuscript, 1974).
18. C. T. C. Wall, *Finiteness conditions for CW complexes*, Ann. of Math. **81** (1965), 56–69.
19. J. E. West, *Mapping Hilbert cube manifolds to ANR's: a solution of a conjecture of Borsuk*, Ann. of Math. **106** (1977), 1–18.

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08544

*Current address:* University of Alabama, Tuscaloosa, Alabama 35487-0350

CALVIN COLLEGE, GRAND RAPIDS, MICHIGAN 49546

*E-mail address:* venema@calvin.edu